

S. Das^a
A.K. Das^b
R.A. Murphy^b
I.C. Punwani^a
M.P. Nasution^c
A.D. Kinghorn^c

^a Department of Pediatric Dentistry,

^b Department of Oral Medicine and
Diagnostic Sciences, College of Dentistry,

^c Department of Medicinal Chemistry and
Pharmacognosy, College of Pharmacy,
University of Illinois at Chicago, Ill., USA

Evaluation of the Cariogenic Potential of the Intense Natural Sweeteners Stevioside and Rebaudioside A

Key Words

Cariogenicity testing
Intense sweeteners
Rebaudioside A
Stevioside
Sugar substitutes

Abstract

Stevioside and rebaudioside A, two intense natural sweeteners, that are constituents of the South American plant *Stevia rebaudiana*, were tested for cariogenicity in albino Sprague-Dawley rats. Sixty rat pups colonized with *Streptococcus sobrinus* were divided into four groups and fed stevioside, rebaudioside A or sucrose added to basal diet 2000 as follows: group 1, 30% sucrose; group 2, 0.5% stevioside; group 3, 0.5% rebaudioside A, and group 4, no addition. All four groups were sacrificed after 5 weeks. *S. sobrinus* counts were made and caries was evaluated according to Keyes' technique. There were no differences in food and water intake and weight gains between the four groups. There were significant differences in sulcal caries scores ($p < 0.02$) and *S. sobrinus* counts ($p < 0.05$) between group 1 and the other three groups. There were no significant differences between the stevioside, rebaudioside A and no-addition groups. It was concluded that neither stevioside nor rebaudioside A is cariogenic under the conditions of this study.

Fermentable carbohydrates are implicated in the etiology of dental caries in many studies. It is also proven that certain oral bacteria, in particular mutans streptococci, ferment these sugars producing acids which in turn cause caries [Rugg-Gunn and Edgar, 1985].

A search for alternatives to sucrose has been in progress for many years, and several intense artificial sweeteners are on the market. A number of natural products of plant origin have been discovered to be potently sweet compounds as a result of a combination of folkloric and laboratory observations [Kinghorn and Soejarto, 1989]. One of the best examples is the sweet plant *Stevia rebaudiana*, which has been used for centuries in Paraguay, its country of origin, to sweeten beverages. Stevioside and re-

baudioside A are the major sweet diterpene glycoside constituents of *S. rebaudiana* leaves. Purified stevioside and extracts of *S. rebaudiana* leaves have been accepted for general use to sweeten a variety of foods and beverages in Japan. Stevioside offers particular advantages over other noncaloric sucrose substitutes in being heat-stable, somewhat resistant to acid hydrolysis, and nonfermentable [Kinghorn and Soejarto, 1991]. Rebaudioside A is present in the leaves of *S. rebaudiana* at lower levels than stevioside, but is a more pleasant-tasting and more water-soluble sweet substance [Crammer and Ikan, 1987]. Both stevioside and rebaudioside A are odorless, white powders. *S. rebaudiana* is cultivated commercially in several Asian countries for the Japanese market. Interest in the poten-

Received:
October 7, 1991
Accepted after revision:
March 22, 1992

Sumitra Das, DDS, MS
Department of Pediatric Dentistry, College of Dentistry
University of Illinois at Chicago
801 South Paulina
Chicago, IL 60612 (USA)

©1992 S. Karger AG, Basel
0008-6568/92/0265-0363
\$ 2.75/0

Table 1. Teklad diet 2000 (cariogenic) catalog No. 170250

	g/kg
Skim milk, powder	280.0
Alfalfa meal (17% protein)	30.0
Desiccated liver (whole liver substance)	10.0
Brewer's yeast, powder	40.0
Wheat flour, whole	60.0
Sucrose, powdered (confectioner's)	560.0
Salt, iodized	20.0

From Keyes and Jordan [1964], modified.

Basal diet 2000 = cariogenic diet 2000 minus sucrose; experimental diet = basal diet 2000 + 56% cornstarch.

tial use of stevioside and rebaudioside A as sucrose substitutes is increasing in the United States and Western Europe [Kingham and Soejarto, 1985, 1991; Phillips, 1987]. Published safety studies on stevioside and extracts of *S. rebaudiana* are quite numerous [Kingham and Soejarto, 1985, 1991]. Of particular importance is an extensive study showing a general lack of chronic toxicity in rodents [Yamada et al., 1985]. However, there are relatively few reports on the effects of stevioside or rebaudioside A related to caries. An in vitro study [Yabu et al., 1977] reported that stevioside added to cultures of *Streptococcus mutans*, *Lactobacillus plantarum* and *Lactobacillus casei* did not support bacterial growth. In another report [Berry and Henry, 1981] it was found that stevioside reduced growth and acid production of *S. mutans* thereby suggesting a cariostatic potential. However, there appear to be no in vivo studies on either sweetener to date. Accordingly, we decided to study the effects of the two sweeteners in an animal (albino rat) model.

Methods and Materials

Test Compounds

The sweet diterpene glycosides stevioside and rebaudioside A were isolated in pure form from the leaves of *Stevia rebaudiana* (Bertoni) Bertoni by gravity-column chromatography, as previously described [Kingham et al., 1982].

Animal Study

The method described below is an accepted procedure [Larson et al., 1977]. We have used this procedure as well as the *S. sobrinus* strain in our previous studies investigating the cariogenicity of aspartame [Das et al., 1982, 1989, 1991].

Pregnant Sprague-Dawley rats were obtained from Charles River Laboratories, Kingston, Mass., USA. They were inoculated with

S. sobrinus 6716-13 resistant to streptomycin (200 mg/ml), according to Larson et al. [1977]. This was done so that the offspring would be infected via coprophagy. All dams received deionized double-distilled water and cariogenic diet 2000 (Teklad). Sixty pups were weaned at 17 days of age. Their oral cavities were checked for the presence of streptomycin-resistant *S. sobrinus* and reinoculated if necessary. All pups received fresh cultures of *S. sobrinus* (2×10^3 cfu/ml) in the drinking water on days 18, 19 and 20. The animals were divided randomly into four groups, so that each group had 15 animals. The experimental animals were fed basal diet 2000 (cariogenic diet 2000 without sucrose) with 56% cornstarch (table 1). Part of this cornstarch was replaced by sucrose, stevioside or rebaudioside A as follows: group 1: 30% sucrose (positive control), group 2: 0.5% stevioside, group 3: 0.5% rebaudioside A and group 4: no replacement (negative control). The 0.5% concentration of stevioside and rebaudioside A was chosen since both are intensely sweet and would accordingly be used in the human diet at very low levels.

The animals were on their respective diets for 5 weeks. During the experiment their diet, water intake and weight were monitored. At the end of the experiments, the animals were sacrificed using a lethal dose of sodium pentothal injected intraperitoneally and then decapitated. Immediately after sacrifice, plaque was collected from all the molars with a sterile swab and used to quantitate *S. sobrinus* in the oral cavities of the rats. The heads were then defleshed, and upper and lower jaws were coded at this time using numbered labels at random from a jar to obscure their group identity. Both upper and lower jaws with molars in place were stained with murexide (0.06% in 70% alcohol) and hemisectioned into buccal and lingual halves. Caries was evaluated according to Keyes' [1958] technique. After scoring the caries, the labels were removed to decode and obtain the animal group and number.

Plaque samples were obtained and quantitatively analyzed. The samples were numbered at random and the group identity was not known to the operator and revealed only at the end. The collected plaque was placed into a vial containing 1 ml of sterile, reduced, anaerobic transport medium of Bacto-Todd Hewitt broth, 0.1% agar and 0.025% resazurin. The vials were then mildly sonicated for three 10-second pulses with 10-second intervals [Madison et al., 1991] after lowering them into the water in a Branson 80-watt Sonic Cleaner. Samples were mixed after dispersion, using a 100- μ l push-button micropipette by repeatedly drawing the fluid into the pipette tip and expelling it back into the vial. A 1:100 dilution of the fluid was prepared using sterile reduced transport medium. Volumes of 0.001, 0.01 and 0.1 ml of the original and the diluted fluid were placed onto the surface of Mitis-Salivarius streptomycin agar plates, using 1-, 10- and 100- μ l push-button micropipettes and spread over the surfaces of the plates using glass spreaders. The plates were incubated anaerobically for 3 days in GasPak jars, and the total number of colony-forming units of *S. sobrinus* was determined from these plates.

All data for food intake, body weight gain, caries incidence and plaque *S. sobrinus* population were analyzed using one-way analysis of variance followed by the Student-Newman-Keuls (SNK) test to identify group means that were statistically different.

Table 2. Caries units [Keyes, 1958] and *S. sobrinus* counts (colony-forming units) in the four groups

	Group 1 30% sucrose	Group 2 0.5% stevioside	Group 3 0.5% rebaudioside A	Group 4 no addition
Animals, n	15	15	15	15
Weight gain, g	162.00 (21.30)	164.80 (13.60)	140.00 (20.80)	150.80 (24.00)
<i>S. sobrinus</i>	200 (10)	90 (5)	91 (8)	88 (9.5)
Sulcal E ¹	15.53 (0.58)	2.75 (0.76)	2.33 (0.60)	1.60 (1.77)
Sulcal Ds ²	4.50 (0.5)	0.50 (0.02)	1.2 (0.08)	0
Smooth surface E	6 (0.60)	0	0	0

Significance levels for intergroup comparisons by Anova and SNK tests: *S. sobrinus*: group 1 vs. 2, 3, 4, $p = 0.05$. Sulcal E caries: group 1 vs. 2, 3, 4, $p = 0.02$. Sulcal Ds caries: group 1 vs. 2, 3, 4, $p = 0.09$. Significance level $p = 0.05$: Differences between groups 2, 3, 4 were not significant ($p > 0.08$). No differences in weight gain between groups. Numbers in parentheses are standard errors.

¹ Enamel caries.

² Slight dentinal caries.

Results

Animals in all four groups appeared to be in good condition throughout the experiment. There were no significant differences in weight gains in the four groups (table 2). The food and water intake was similar in all groups.

S. sobrinus counts: The control group (group 1) fed 30% sucrose in diet had the highest number (statistically significant) of *S. sobrinus* among the four groups ($p = 0.05$). There were no significant differences in *S. sobrinus* counts between the remaining three groups (table 2).

Sulcal caries scores were significantly higher ($p = 0.02$) in the animals fed sucrose. No significant differences were found between the other three groups which had very low incidences of sulcal caries (table 2). Dentinal (sulcal) caries (Ds) was slightly higher (but statistically insignificant, $p = 0.10$) in the sucrose-fed animals than the two experimental groups. The animals receiving no additive had no dentinal caries. Smooth surface caries was found only in the group receiving sucrose.

Discussion

Neither stevioside nor rebaudioside A produced any adverse reactions in the animals. This was not surprising, considering previous reports of the lack of toxicity of *S. rebaudiana* plant and stevioside in animal models [Kingham and Soejarto, 1985; Yamada et al., 1985] and the present widespread consumption of *S. rebaudiana* products by humans in Japan [Kingham and Soejarto, 1991].

The *S. sobrinus* populations in the experimental groups receiving either stevioside or rebaudioside A were comparable to that in group 4, which was not supplied with any additive. This suggests that under the conditions of this study, stevioside and rebaudioside A had no detectable effect on the colonization and growth of *S. sobrinus* in the oral cavities of the experimental animals. This is consistent with the in vitro findings of Yabu et al. [1977], who reported that stevioside, when added to cultures of *S. mutans*, *L. plantarium* and *L. casei*, did not support their growth, and with those of Berry and Henry [1981], who found that stevioside-supplemented medium reduced the growth of *S. mutans* in comparison to the medium with sucrose. It can be argued that the concentration of the sweeteners used in this experiment was very low in comparison to that of sucrose. However, due to the fact that both these intense sweeteners are 300 or more times sweeter than sucrose, their normal human usage would be at very low concentrations.

The incidence of caries was directly correlated with the presence of sucrose and the concomitant increase in *S. sobrinus*. Sulcal caries was high in sucrose-fed animals as was the *S. sobrinus* count, and this was the only group with smooth-surface caries.

We have not studied the effects of either stevioside or rebaudioside A on caries production in the presence of sucrose. A previous in vitro study by Yabu et al. [1977] suggested a cariostatic potential of stevioside when added to a sucrose medium by suppressing bacterial growth. Similarly, stevioside lowered acid production by *S. mutans* in a sucrose medium [Berry and Henry, 1981]. Studying the ef-

fects of these intense sweeteners on caries in the presence of sucrose should be the next phase in future experimental research. However, in the present experiment, we have found the two sweeteners to be noncariogenic.

Acknowledgments

We are grateful to Shirley Thomas and Aretha Draper for their clerical assistance.

This work was supported by Wach Fund award No. 2-6-46507 from the University of Illinois.

References

- Berry CW, Henry CA: Effect of stevioside on the growth and acid production of *Streptococcus mutans*. J Dent Res 1981;60:479.
- Crammer B, Ikan R: Progress in the chemistry and properties of rebaudioside; in Greenby TH (ed): Developments in Sweeteners. London, Elsevier, 1987, vol 3, pp 45-64.
- Das S, Das AK, Mukherjee S: Effect of molybdenum on dental caries incidence in rats. J Dent Res 1982;61:327.
- Das S, Das AK, Murphy RA: Evidence for anticariogenicity of aspartame. J Dent Res 1989;68:246.
- Das S, Das AK, Murphy RA, Varawangvasu R: Aspartame and dental caries in the rat. Pediatr Dent 1991;13:217-220.
- Keyes PH: Dental caries in the molar tooth of rats. A method for diagnosing and scoring several types of lesions simultaneously. J Dent Res 1958;37:1088-1099.
- Keyes PH, Jordan HV: Periodontal lesions in the Syrian hamster. III. Findings related to an infectious and transmissible component. Arch Oral Biol 1964;9:337-440.
- Kinghorn AD, Nanayakkara NPD, Soejarto DD, Medon PJ, Kamath S: Potential sweetening agents of plant origin. I. Purification of *Stevia rebaudiana* sweet constituents by droplet counter-current chromatography. J Chromatogr 1982;237:478-482.
- Kinghorn AD, Soejarto DD: Current status of stevioside as a sweetening agent for human use; in Wagner H, Hikino H, Farnsworth NR (eds): Economic and Medicinal Plant Research. London, Academic Press, 1985, vol 1.
- Kinghorn AD, Soejarto DD: Intensely sweet compounds of natural origin. Med Res Rev 1989;9:91-115.
- Kinghorn AD, Soejarto DD: Stevioside; in O'Brien Nabors L, Gelardi RC (eds): Alternative Sweeteners. New York, Dekker, 1991, vol 2, pp 157-171.
- Larson RH, Almsbaugh SM, Navia JM, Rosen S, Schuster GS, Shaw JH: Collaborative evaluation of rat caries model in six laboratories. J Dent Res 1977;56:1007-1012.
- Madison KM, Bowen WM, Pearson SK, Falany JL: Enhancing the virulence of *Streptococcus sobrinus* in rats. J Dent Res 1991;70:38-43.
- Phillips KC: Stevia: Steps in developing a new sweetener; in Greenby TH (ed): Developments in Sweeteners. London, Elsevier, 1987, vol 3, pp 1-43.
- Rugg-Gunn AJ, Edgar WM: Sweeteners and dental health. Community Dent Health 1985;2:213-223.
- Yabu M, Takase M, Toda K, Tanimoto K, Yasutake A, Iwamoto Y: Studies on stevioside natural sweetener: Effect on some oral organisms. Hiroshima Daigaku Shigaku Zasshi 1977;9:12-17.
- Yamada A, Ohgaki S, Noda T, Shimizu M: Chronic toxicity study of dietary Stevia extracts in F344 rats. J Food Hyg Soc 1985;26:169-183.